

Spatio-Temporal Patterns for a Generalized Innovation Diffusion Model

Fariba Hashemi

Ecole Polytechnique Fédérale de Lausanne (EPFL), College of Management of Technology (CDM),
Management of Technology and Entrepreneurship Institute (MTEI)

Max-Olivier Hongler

Ecole Polytechnique Fédérale de Lausanne (EPFL), School of Engineering (STI),
Laboratory of Microengineering for Manufacturing (LPM)

Olivier Gallay

Ecole Polytechnique Fédérale de Lausanne (EPFL), School of Architecture, Civil and Environmental
Engineering (ENAC), Transport and Mobility Laboratory (Transp-OR)

August 31, 2010

Abstract

We construct a model of innovation diffusion that incorporates a spatial component into a classical imitation-innovation dynamics first introduced by F. Bass. Relevant for situations where the imitation process explicitly depends on the spatial proximity between agents, the resulting nonlinear field dynamics is exactly solvable. As expected for nonlinear collective dynamics, the imitation mechanism generates spatio-temporal patterns, possessing here the remarkable feature that they can be explicitly and analytically discussed. The simplicity of the model, its intimate connection with the original Bass' modeling framework and the exact transient solutions offer a rather unique theoretical stylized framework to describe how innovation jointly develops in space and time.

Keywords: Diffusion of innovation - Bass' model - interactive multi-agent systems - local interactions - imitation processes - stochastic dynamics - alternating renewal processes - mean-field dynamics - discrete velocity collisions models - nonlinear field equations - exact transient evolutions.

1 Introduction

Since the middle of the XXth century, a substantial literature emphasizes the importance of quantitative models that enable the forecasting of the diffusion of technological innovations (DTI) not only for pure academic interest, but for its practical relevance. While the empirical evidence that quantification of DTI is possible was initially recognized by E. Mansfield [1] and Z. V. Griliches [2, 3], the first quantitative stylized dynamical modeling framework was proposed by F. Bass in his seminal 1969 paper [4]. Similar in essence to the P.-F. Verhulst's epidemiological logistic equation, the Bass' model uses an **aggregated differential approach** enabling the reproduction of the relevant dynamical features of the adoption of a new product and/or a new technology in a society of consumers. Bass' nonlinear dynamics is basically governed by the ratio of two control parameters, namely the *innovation* and *imitation* rates. Introducing a quadratic nonlinearity into the evolution equation, the model mathematically describes consumer imitative interactions. This nonlinearity leads to an evolution characterized by two distinct time scales, *i.e.* a fast initial exponentially growing phase followed by a slow asymptotic evolution when full equilibrium demand is nearly reached. The seminal works of J. A. Schumpeter [5] and subsequently of B. Jovanovic and R. Rob [6] support this observation by illustrating the importance of imitation waves in DTI and the formation of business cycles.

In general, interaction-based models have been employed for various applications ranging from epidemiology [7], variance of crime rates across space when direct interdependencies occur between nearest neighbors [8], herd behavior in financial markets [9, 10, 11, 12] to social movements and political uprisings (see [13] and [14] for several additional references). Interaction-based methods have also been useful tools in modeling the diffusion of innovation [16, 17, 18, 19, 20, 21]. In their contribution, G. Ellison and D. Fudenberg [22, 23] study agents who consider the experiences of their neighbors in deciding which of two technologies to adopt, in a world where players use rules of thumb that ignore historical data but may incorporate a tendency to use the more popular technology. In this learning environment, agents ob-

serve both their neighbors' choices, and periodically reevaluate their decisions, as opposed to making a once-and-for-all choice. R. Andergassen et al. [24] investigate the evolutionary process of imitation and innovation as a search mechanism in a given neighbourhood of firms. In their world, the spreading of information through neighbourhoods allows firms to acquire knowledge leading to innovation waves as firms attempt to glean information on best practice techniques, which they subsequently imitate. For additional models where preference orderings over alternatives in a choice set can depend on the actions chosen by other agents, see [25, 26, 27, 28].

The original Bass' dynamics is aggregated and hence fails **to detail the influence of spatial location of agents** in the imitative behavior of consumers. Intuitively, spatial considerations strongly affect the interactions between agents and the underlying imitation mechanisms. In fact spatial proximity is argued to be a major driving force of innovation diffusion which often exceeds the external marketing efforts such as advertising [29, 30]. In 1991, P. Krugman [31] pointed out that production is remarkably concentrated in space. This observation opened a strong research effort devoted to the understanding of the spatial dimension of innovation diffusion. M. P. Feldman [32, 33] echoed P. Krugman's observation in pointing out that geographical effects are even more stringent for innovative activity because the rationale for the formation of adopter clusters is related to the role of word-of-mouth and imitation in the diffusion of innovations. As emphasized in [34], a clear correlation exists between geographical proximity and the strength and speed of word-of-mouth spread, sometimes labeled as the *neighborhood effect*.

In addition to these works, a wealth of empirical studies exemplify the importance of geography in the diffusion of knowledge and R & D. Spatially-mediated knowledge spillovers of R & D are explicitly discussed in [35, 36, 37, 38]. It is noteworthy to observe that this pure geographic view can be generalized by defining metric distances on abstract state spaces in order to describe the evolution of technological advance, R & D investment volume or any other abstract features [39] on which agents can compete by adjustment of their individual behavior.

Recent literature suggests that imitation interactions between interacting agents like bacteria, flies, quadrupeds or fishes can explain the formation of compact spatio-temporal patterns, *i.e.* *swarms* or *platoons*, which spatially evolve as quasi-solid bodies [40, 41]. The *flock-*

ing mechanism originates from mimetic type decisions based on agents' observations of their neighbors. To the best of our knowledge, spatial flocking mechanisms seem to be barely discussed in the interaction-based socioeconomic literature. Hence a natural and simple attempt to analytically infer the role of spatial parting is to introduce spatial effects and it is the aim of our paper to incorporate their influence into the original Bass' evolution.

Adding a spatial dimension transforms the Bass' ordinary differential equation into a partial differential equation (PDE). Due to the underlying imitation mechanism, the resulting PDE will be intrinsically nonlinear, a perspective that generally offers little hope for explicit solutions in the realm of field theories. The present paper illustrates how a simple natural spatial extension of the original Bass' dynamics leads nevertheless to a fully solvable nonlinear field dynamics, a truly remarkable result. The resulting equations belong to the discrete velocities Boltzmann equations (DVBE) which describe the macroscopic properties of a dilute gas. The specific DVBE that can be derived from the Bass' dynamics coincide with the Ruijgrok-Wu (RW) model introduced and solved by T. W. Ruijgrok and T. T. Wu [42]. This intimate connection with statistical physics suggests that the Bass' dynamics can be obtained, via a *mean-field* limit, from a microscopic point of view in which a large number of agents interact. While the mean-field approach is a basic tool in statistical physics of large systems, it has now been explicitly used in recent econometric studies as well (see illustrations in [10, 11, 12, 14, 15, 28]). Relying on the law of large numbers, the mean-field limit allows one to write deterministic evolution for probability densities in question. In the sequel, we will explicitly construct the microscopic connection that exists between the Bass' imitation model with spatial effects and the RW model inspired by a similar approach adopted in [43] for a multi-agent dynamics in logistics and econophysics contexts. In [43], the dynamics exhibit a nonlinear term due to a specific imitation mechanism giving rise to the famous Burgers' nonlinear PDE to describe the emergence of spatio-temporal patterns. As illustrated in [44], RW dynamics actually generalizes the Burgers' equation, the spatio-temporal Bass' model presented here can itself be viewed as a natural generalization of the multi-agent imitation model studied in [43].

Besides its direct practical relevance, the simplicity of the original Bass model, for which exact analytical solutions are available, has undoubtedly contributed to its popularity in the economics and management literatures. Endowing Bass' dynamics with spatially-dependent imita-

tion mechanisms confers a new dimension to interaction-based socio-economic modeling, opening the possibility to analytically study the generation of spatio-temporal patterns in a highly nonlinear context. Our stylized dynamics can be viewed as an exceptional possibility to analytically observe the spatio-temporal effects arising for a collection of agents subject to imitation interactions.

2 Spatially-Dependent Imitation Dynamics

Consider a collection \mathcal{A} of N autonomous agents which are in a migration process on the one-dimensional real line \mathbb{R} . At any time $t \in \mathbb{R}^+$, we assume that the complete population is composed by two types of agents \mathcal{A}_+ and \mathcal{A}_- , ($\mathcal{A} = \mathcal{A}_- \cup \mathcal{A}_+$), characterized by two associated (x, t) -dependent migration velocities $V_+(x, t)$ and $V_-(x, t)$ on \mathbb{R} . At any time, each agent is free to modify his/her velocity. Agents $a_k \in \mathcal{A}$, $k = 1, 2, \dots, N$, flip their velocities either spontaneously or after an autonomous decision based on an imitation process (IP). Let us write $\alpha(x, t)$, respectively $\beta(x, t)$, as the spontaneous transformation rates from states $\mathcal{A}_+ \mapsto_{\alpha(x,t)} \mathcal{A}_-$, respectively $\mathcal{A}_- \mapsto_{\beta(x,t)} \mathcal{A}_+$. Apart from these spontaneous transitions, additional transitions are assumed to be triggered by mutual agents' interactions. Specifically, for agent a_k , the IP mechanism is assumed to depend on the observation of the present velocity states, (*i.e.* V_+ or V_-), adopted by other proximity members located in the neighborhood $\mathcal{N}_k = \mathcal{N}_{k,-} \cup \mathcal{N}_{k,+} \subset \mathcal{A}$ of agent a_k . For an arbitrary agent $a_{k,\pm}$, we define his/her imitation decision rule according to his/her interactions with other agents as follows:

i) Dynamic rule for an agent $a_{k,-}$. At time t , the agent $a_{k,-}$ simultaneously observes the (velocity) states of the agents contained in his/her neighborhood \mathcal{N}_k . The presence of agents $a_{j,+} \in \mathcal{N}_k$, $j \neq k$, seen by $a_{k,-}$ triggers an imitation mechanism which enhances the transition rate towards state V_+ , *i.e.* $\beta(x, t) \mapsto [\beta(x, t) + i_k(x, t)]$. The extra contribution $i_k(x, t)$ is proportional to (monotonically increasing with) the number N_k of agents $a_{j,+}$ in the neighborhood \mathcal{N}_k of agent $a_{j,-}$ (*i.e.* $N_k = \text{card}(\mathcal{N}_k)$).

ii) Dynamic rule for an agent $a_{k,+}$. At time t , the agent $a_{k,+}$ simultaneously observes the (velocity) states of the agents contained in his/her neighborhood \mathcal{N}_k and **does not modify its velocity** whatever he/she observes.

According to these dynamic rules, the time-dependent position $X_k(t)$ of agent a_k , $k = 1, 2, \dots, N$, can be written as a set of coupled stochastic differential equations (SDEs):

$$\dot{X}_k(t) = I_k(t), \quad k = 1, 2, \dots, N, \quad (1)$$

where $I_k(t)$ stands for a two-velocity-states Markov chain (*i.e.* the state space is here $\Omega := \{V_-(x, t), V_+(x, t)\}$) the transition rates of which are defined by :

$$V_+(x, t) \mapsto_{\alpha(x, t)} V_-(x, t) \quad V_-(x, t) \mapsto_{[\beta(x, t) + i_k(x, t)]} V_+(x, t). \quad (2)$$

The noise source in Eq.(1) can also be viewed as a non-homogeneous, alternating Markov renewal process in which the inverse transition rates $\alpha(x, t)^{-1}$ and $[\beta(x, t) + i_k(x, t)]^{-1}$ are respectively the mean sojourn times in states $V_+(x, t)$ and $V_-(x, t)$. Observe therefore that in Eq.(1), the coupling between the various agents is realized via the extra $i_k(x, t)$ transition rate.

In what follows, we consider a very large population of agents, *i.e.* $N \rightarrow \infty$, and we focus on observation neighborhoods \mathcal{N}_k , $k = 1, 2, \dots, N$, that contain on average $N_k = \rho N$ agents ($0 < \rho < 1$), implying in turn that $N_k \rightarrow \infty$. Instead of individual trajectories, let us now think in terms of statistical properties characterizing the evolution of the whole population of agents \mathcal{A} . To this aim, we write $P(x, t) \in [0, 1]$ and $Q(x, t) \in [0, 1]$ to denote the density of agents \mathcal{A}_+ and \mathcal{A}_- to be found at position $x \in \mathbb{R}$ at time t .

We now adopt a **mean-field approach (MFA)**, which consists in considering that, statistically, the time evolution of an arbitrary agent, say $a_k \in \mathcal{A}$, is representative of the whole population and the influence due to his/her neighbors located in \mathcal{N}_k is viewed as an **external effective interactive mean field**. Accordingly, the MFA allows us to replace Eq.(1) by a single scalar SDE:

$$\dot{X}(t) = I(t), \quad (3)$$

where in Eq.(3), the $i_k(x, t)$ transition rates are replaced by an effective rate $i(x, t)$ reading as:

$$i(x, t) = \lim_{\mathcal{N} \rightarrow \infty} \frac{1}{\mathcal{N}} \sum_{j=1}^{\mathcal{N}} a_{j,+} = \int_x^{x+\Gamma} P(|x-z|, t) dz. \quad (4)$$

with $\Gamma \in \mathbb{R}^+$ characterizing the typical neighborhood interval \mathcal{N} .

Note that while we proceeded here heuristically, a rigorous mathematical justification for the validity of the MFA for this type of dynamics relies on the property of *propagation of chaos*, as explained in [47] for SDEs of the type given by Eq.(1) when the noise is a White Gaussian Noise (WGN). SDEs of the type given by Eq.(1) driven by WGN have recently been used for multi-agent dynamics with imitation process in [43].

Associated with the SDEs given by Eqs.(3) and (4), the MFA enables us to write a Fokker-Planck equation for the probability densities $P(x, t)$ and $Q(x, t)$, [44]:

$$\begin{aligned} \dot{P}(x, t) + V_+(x, t) \frac{\partial}{\partial x} P(x, t) &= +\mathcal{J}(P(x, t), Q(x, t)) - \alpha P(x, t) + \beta Q(x, t), \\ \dot{Q}(x, t) + V_-(x, t) \frac{\partial}{\partial x} Q(x, t) &= -\mathcal{J}(P(x, t), Q(x, t)) + \alpha P(x, t) - \beta Q(x, t), \end{aligned} \quad (5)$$

where the **imitation rate term** in Eqs.(5) reads as:

$$\mathcal{J}(P(x, t), Q(x, t)) = Q(x, t) \left[\int_x^{x+\Gamma} P(|x-z|, t) dz \right]. \quad (6)$$

The dynamics in Eqs.(5) is a coupled set of **nonlocal and nonlinear field equations**, which barely offers hope for any analytical discussion. However, for small Γ (*i.e.* infinitesimal interaction neighborhoods), we may Taylor expand Eqs.(5), up to first order in Γ , to obtain:

$$\begin{aligned} \dot{P}(x, t) + V_+(x, t) \frac{\partial}{\partial x} P(x, t) &= +\Gamma P(x, t) Q(x, t) - \alpha P(x, t) + \beta Q(x, t), \\ \dot{Q}(x, t) + V_-(x, t) \frac{\partial}{\partial x} Q(x, t) &= -\Gamma P(x, t) Q(x, t) + \alpha P(x, t) - \beta Q(x, t). \end{aligned} \quad (7)$$

We directly observe that Eqs.(7) can be viewed as a generalized Bass' dynamics which confers relevance to the spatial dimension on which agents evolve. Indeed, similarly to the original Bass' model, we include an imitation process, represented in Eqs.(7) by the nonlinear contribution $\Gamma P(x, t) Q(x, t)$.

Writing $P(x, t) + Q(x, t) = \Sigma(x, t)$, the summation of both equations in (7) yields a continuity equation:

$$\frac{\partial}{\partial t} \Sigma(x, t) + \frac{\partial}{\partial x} \left[\left(\frac{V_+(x, t) + V_-(x, t)}{2} \right) \Sigma(x, t) \right] = 0, \quad (8)$$

which in turn implies the normalization constraint:

$$\int_{\mathbb{R}} P(x, t) dx + \int_{\mathbb{R}} Q(x, t) dx \equiv 1, \quad \forall t \in \mathbb{R}^+. \quad (9)$$

Although the dynamics given by Eqs.(7) has been derived for non-homogeneous and non-stationary parameters $\alpha(x, t)$, $\beta(x, t)$ and $V_{\pm}(x, t)$, in the sequel we will restrict our attention to situations where these parameters can be assimilated to constants (*i.e.* α, β and V_{\pm}).

2.1 Spatial Homogeneous Regimes - Bass' Model

When $V_- = V_+ = 0$, the spatial character disappears from the dynamics given by Eqs.(7), which implies that $P(x, t) \equiv P(t)$ and $(Q(x, t) = Q(t)$. More precisely, Eqs.(7) becomes:

$$\begin{aligned} \dot{P}(t) &= +\Gamma P(t)Q(t) - \alpha P(t) + \beta Q(t), \\ \dot{Q}(t) &= -\Gamma P(t)Q(t) + \alpha P(t) - \beta Q(t), \end{aligned} \quad (10)$$

with the notation $\dot{P}(t) := \frac{d}{dt}P(t)$. The constraint given by Eq.(9) now simply becomes $P(t) + Q(t) = 1$ and enables us to rewrite Eq.(10) in the following form:

$$\dot{P}(t) = -\Gamma \left\{ P^2(t) - \left[1 - \frac{(\alpha + \beta)}{\Gamma} \right] P(t) - \frac{\beta}{\Gamma} \right\}. \quad (11)$$

At this stage, it is worth observing that for $\alpha = 0$ and $\Gamma = 1$ (without loss of generality, indeed it can always be obtained by the time rescaling $t \mapsto \tau = \Gamma t$), Eq.(11) reduces to:

$$\dot{P}(t) = [1 - P(t)] [\beta + P(t)], \quad (12)$$

which is precisely the original Bass' dynamics [4] with β being the ratio between the *imitation* and *innovation* rates.

For completeness of the exposition, let us integrate Eq.(11), with the initial condition $P(t = 0) = P_0$, to get:

$$P(t) = \frac{(\Delta + b)(P_0 - \Delta + b)e^{-2\Delta\Gamma t} + (\Delta - b)(P_0 + \Delta + b)}{(P_0 + \Delta + b) - (P_0 - \Delta + b)e^{-2\Delta\Gamma t}} \quad (13)$$

with the definitions:

$$b = -\frac{1}{2\Gamma}(\Gamma - \alpha - \beta) \quad \text{and} \quad \Delta = \frac{1}{2\Gamma}\sqrt{(\Gamma - \alpha - \beta)^2 + 4\beta}.$$

In the asymptotic time limit $t \rightarrow \infty$, Eq.(13) converges to:

$$\lim_{t \rightarrow \infty} P(t) = P_{\text{station}} = \Delta - b. \quad (14)$$

For the original Bass' model obtained when $\alpha = 0$ and $\Gamma = 1$, we have $b = -\frac{1}{2}(1 - \beta)$ and $\Delta = \frac{1}{2}(1 + \beta)$. Accordingly, with $P_0 = 0$, Eq.(13) reduces to the original Bass' solution:

$$P(t) = \frac{\beta [1 - e^{-(1+\beta)t}]}{\beta + e^{-(1+\beta)t}}, \quad Q(t) = \frac{(1 + \beta)e^{-(1+\beta)t}}{\beta + e^{-(1+\beta)t}}. \quad (15)$$

Moreover $P_{\text{station}} = \Delta - b = 1$, which expresses the fact that all agents ultimately adopt the new technology, as it is obviously expected in the original Bass' modeling framework, when $\alpha = 0$.

3 Bass' Dynamics with Spatio-Temporal Effects

Coming back to the general dynamics given by Eqs.(7) and introducing dimensionless coordinates via the Gallileo transformation $(x, t) \mapsto (y, s)$ defined by:

$$\begin{pmatrix} y \\ s \end{pmatrix} = \begin{pmatrix} \frac{2\Gamma}{(V_+ - V_-)} & \frac{\Gamma(V_- + V_+)}{(V_- - V_+)} \\ 0 & \Gamma \end{pmatrix} \begin{pmatrix} x \\ t \end{pmatrix}, \quad (16)$$

we straightforwardly have:

$$\partial_x(\cdot) \mapsto \left[\frac{2\Gamma}{(V_+ - V_-)} \right] \partial_y(\cdot) \quad \text{and} \quad \partial_t(\cdot) \mapsto \left[\frac{\Gamma(V_- + V_+)}{(V_- - V_+)} \right] \partial_y(\cdot) + \Gamma \partial_s(\cdot), \quad (17)$$

which transforms Eqs.(7) into the coupled set of nonlinear partial differential equations (PDEs):

$$\begin{aligned} \partial_s P + \partial_y P &= +PQ - \frac{\alpha}{\Gamma} P + \frac{\beta}{\Gamma} Q, \\ \partial_s Q - \partial_y Q &= -PQ + \frac{\alpha}{\Gamma} P - \frac{\beta}{\Gamma} Q. \end{aligned} \quad (18)$$

The set of non-linear PDEs given by Eqs.(18) can be interpreted as being a discrete two velocity model of Boltzmann equations, first studied in [42]. The dynamics given by Eqs.(18) is remarkable, as using a generalized Hopf-Cole logarithmic transformation, we get:

$$P(y, s) = -\frac{\beta}{\Gamma} + \partial_s \log H(y, s) - \partial_y \log H(y, s) \quad (19)$$

$$Q(y, s) = \frac{\alpha}{\Gamma} - \partial_s \log H(y, s) - \partial_y \log H(y, s) \quad (20)$$

which actually reduces the set Eqs.(18) into the *Telegraphist equation*:

$$\partial_{ss}H(y, s) - \partial_{yy}H(y, s) - \frac{\alpha\beta}{\Gamma^2}H(y, s) = 0. \quad (21)$$

Accordingly, the dynamics given by Eqs.(7) with constant parameters has the **truly exceptional property that it can be exactly and explicitly solved for any initial conditions** $P_0(y)$ and $Q_0(y)$. According to [42], the general solution reads as:

$$H(y, s) = \frac{1}{2} [A(y+s) + A(y-s)] + \frac{1}{2}\mathcal{B}_1(y, s) + \frac{\eta s}{2}\mathcal{B}_2(y, s), \quad (22)$$

where $\eta = \frac{\sqrt{\alpha\beta}}{\Gamma}$ and where we have the following definitions:

$$\mathcal{B}_1(y, s) = \int_{y-s}^{y+s} \mathbb{I}_0 \left(\eta \sqrt{s^2 - (y-y')^2} \right) B(y') dy',$$

and

$$\mathcal{B}_2(y, s) = \int_{y-s}^{y+s} \left(\frac{1}{\sqrt{s^2 - (y-y')^2}} \right) \mathbb{I}_1 \left(\eta \sqrt{s^2 - (y-y')^2} \right) A(y') dy',$$

with $\mathbb{I}_0(\cdot)$ and $\mathbb{I}_1(\cdot)$ being the modified Bessel's functions and:

$$B(y) = \frac{1}{2} \left[P_0(y) - Q_0(y) + \frac{\alpha}{\Gamma} + \frac{\beta}{\Gamma} \right] A(y), \quad (23)$$

$$A(y) = \exp \left\{ -\frac{1}{2} \int^y \left[P_0(y') + Q_0(y') - \frac{\alpha}{\Gamma} + \frac{\beta}{\Gamma} \right] dy' \right\}. \quad (24)$$

3.1 Behavior of the Solutions

Though fully explicit and exact, the truly beautiful solution given by Eqs.(19) and (20) deserves discussion and interpretation for specific situations and this is precisely the objective of this section. In what follows, we will systematically choose $\Gamma = 1$ and unit velocities $V_{\pm} = \pm 1$.

Let us here focus on the Bass' dynamics that results when $\alpha = 0$. This directly implies that $\eta = \sqrt{\alpha\beta} = 0$ and hence Eqs.(19) and (20) coincide with the ordinary wave equation. By definition of Bessel's

functions [48], we have that $\mathbb{I}_0(0) = 1$ and $\mathbb{I}_1(0) = 0$, thus leading to the usual wave solution in the form:

$$H(y, s) = \frac{1}{2} [A(y + s) + A(y - s)] + \frac{1}{2} \int_{y-s}^{y+s} B(y') dy'. \quad (25)$$

Due to the fact that $\alpha = 0$, we expect that the agents' population \mathcal{A}_+ with velocity $V_+ = +1$ increases by opposition to the population \mathcal{A}_- of agents with velocity $V_- = -1$ which is doomed to extinction. Let us now explore the transient nature of the solution and this for three types of initial conditions.

a) Initial conditions: $P_0(\mathbf{y}) = \delta(\mathbf{y})$ and $Q_0(\mathbf{y}) = 0$.

In this case, the time-dependent solution reads as:

$$P(y, s) = \delta(y - s) \quad \text{and} \quad Q(y, s) \equiv 0. \quad (26)$$

as it can be checked directly from Eqs.(19) and (20). Eqs.(26) are clearly consistent with the fact that $\alpha = 0$ and hence that no transitions from $V_+ = +1$ to $V_- = -1$ velocities occur. Hence, starting with all agents with velocity $V_+ = +1$, they stay with their original velocity and the density $P(y, s)$ is a uniformly traveling Dirac mass with velocity $V_+ = +1$ towards the positive \mathbb{R} -axis.

b) Initial conditions: $P_0(\mathbf{y}) = 0$ and $Q_0(\mathbf{y}) = \delta(\mathbf{y})$.

Either by direct substitution into Eqs.(18) or alternatively by using Eqs.(19) and (20), one can verify that the time-dependent solution in this case reads as:

$$P(y, s) = \frac{\beta}{2} e^{\frac{\beta}{2}(y-s)} \Theta(|y|-s) \quad \text{and} \quad Q(y, s) = e^{-\beta s} \delta(y+s) \quad (27)$$

where $\Theta(|y| - s)$ is a Heaviside cutoff function which identically vanishes for negative arguments. The dynamics $P(y, s)$ of the spatial dispersion of the agents with velocity $V_+ = +1$ is illustrated in Fig. 1.

The behavior of the solution given by Eqs.(27) can be easily understood. Indeed, the Dirac mass for $Q(y, s)$ expresses the fact that agents with velocity $V_- = -1$ are gradually depopulated at the rate $s\beta$ for the benefit of agents traveling with velocity $V_+ = +1$ and hence migrating to $P(y, s)$. The Heaviside function




Figure 1: Spatial dispersion $P(y, s)$ of the agents with velocity $V_+ = +1$ for different times $s = [0.01; 20; 40]$. The initial conditions are given by $P_0(y) = 0$ and $Q_0(y) = \delta(y)$, $\alpha = 0$ and $\beta = 0.1$. In this case, the agents are immediately segregated and there are no imitation processes.

$\Theta(|y| - s)$ expresses the fact that no agent can possibly be found at a distance larger than $|y|$ at time t (remember that the velocities here are $V_{\pm} = \pm 1$). It is worth observing that in the original Bass' model, the adoption rate given in Eq.(15) is equal to $(1+\beta)$. This overcomes the β adoption rate that we will obtain below in Eqs.(28), due to the fact that in the present configuration, imitation process does not enter into play in our spatial model, which drastically moderates the overall adoption rate. Indeed, in this limiting regime the agents' populations \mathcal{A}_+ and \mathcal{A}_- are immediately and definitively segregated, which hence never allows imitation processes to take effect (the unit rate discrepancy between the $(1+\beta)$ adoption rate occurring in Eq.(15) and the β rate in Eq.(27) is obtained from the $\Gamma = 1$ choice). The resulting temporal evolution of the agents' overall adoption rate obtained in the present case ($P(t) = \int_{\mathbb{R}} P(y, t) dy$), compared with the one of the original Bass' model, will be illustrated later in Fig. 4.

Finally, let us observe that, from Eqs. (27), one immediately

obtains:

$$\int_{\mathbb{R}} P(y, s) dy = [1 - e^{-\beta s}] \quad \text{and} \quad \int_{\mathbb{R}} Q(y, s) dy = e^{-\beta s}, \quad (28)$$

thus showing that Eq.(9) is fulfilled. In addition, for asymptotic times $s \rightarrow \infty$, Eqs.(28) indicate that all agents ultimately adopt the $V_+ = +1$ velocity as it is expected for the $\alpha = 0$ regime.

c) Initial conditions: $P_0(y) = Q_0(y) = \frac{1}{8\gamma} [\tanh(y + \gamma) - \tanh(y - \gamma)]$.

For these initial conditions and for the particular choice $\beta = 2$ and $\gamma = \frac{1}{4}$, the resulting time-dependent solution of our spatial Bass' model is given by:

$$P(y, s) = \frac{1}{2} \left\{ -2 + \frac{1}{H(y, s)} [A(y - s) + e^{-y+s} \frac{\cosh(y - s - \gamma)}{\cosh(y - s + \gamma)} (1 - \tanh(y - s - \gamma) + \tanh(y - s + \gamma))] \right\} \quad (29)$$

and

$$Q(y, s) = \frac{1}{2} \left\{ \frac{1}{H(y, s)} [-A(y + s) + e^{-y-s} \frac{\cosh(y + s - \gamma)}{\cosh(y + s + \gamma)} (1 - \tanh(y + s - \gamma) + \tanh(y + s + \gamma))] \right\}, \quad (30)$$

where

$$H(y, s) = \frac{1}{2} [A(y + s) + A(y - s) + \mathcal{B}_1(y, s)],$$

$$A(y) = e^y \frac{\cosh(y - \gamma)}{\cosh(y + \gamma)},$$

$$\mathcal{B}_1(y, s) = \frac{e^{-\gamma}}{2} [\text{atan}(\sinh(y + s + \gamma)) - \text{atan}(\sinh(y - s + \gamma))] - e^{3\gamma} [e^{-y-s-\gamma} + \text{atan}(e^{y+s+\gamma}) - e^{-y+s-\gamma} - \text{atan}(e^{y-s+\gamma})].$$

The spatio-temporal dynamics $P(y, s)$ of the agents with velocity $V_+ = +1$ is illustrated in Fig. 2. The joint evolutions of $P(y, s)$ and $Q(y, s)$ (representing agents with velocity $V_+ = +1$ and $V_- = -1$ respectively) are drawn in Fig. 3.

In the present configuration, the two types of agents have an




Figure 2: Spatial dispersion $P(y, s)$ of the agents with velocity $V_+ = +1$ for different times $s = [0; 0.2; 3; 10]$. The initial conditions are given by $P_0(y) = Q_0(y) = \frac{1}{8\gamma} [\tanh(y + \gamma) - \tanh(y - \gamma)]$, $\alpha = 0$, $\beta = 2$ and $\gamma = \frac{1}{4}$.

identical initial spatial distribution (*i.e.* $P_0(y) = Q_0(y)$, $\forall y$). Hence, half of the agents have initially the velocity $V_+ = +1$, the other half having the velocity $V_- = -1$ (*i.e.* $\int_{\mathbb{R}} P_0(y) dy = \int_{\mathbb{R}} Q_0(y) dy = \frac{1}{2}$). The spatio-temporal behavior of the solution given by Eqs.(29) and (30) can be split into two different time phases. For short times of the dynamics, the overlap between $P(y, s)$ and $Q(y, s)$ is non-null (*i.e.* the two populations of agents \mathcal{A}_+ and \mathcal{A}_- are not spatially segregated), thus leading to strong imitation processes between the agents. Agents are changing their velocity from $V_- = -1$ to $V_+ = +1$ due to both imitation and spontaneous transitions. Accordingly, $Q(y, s)$ is gradually depopulated for the benefit of $P(y, s)$. The imitation processes are decreasing as the overlap between $P(y, s)$ and $Q(y, s)$ gets smaller, but the rate β of spontaneous transitions from $V_- = -1$ to $V_+ = +1$ remains itself constant with time. In the time asymptotic regime, almost all the agents have adopted velocity $V_+ = +1$, leading $P(y, s)$ to behave as a probability density uniformly traveling with velocity $V_+ = +1$ towards the positive \mathbb{R} -axis.




Figure 3: Spatial dispersions $P(y, s)$ and $Q(y, s)$ of the agents with velocity $V_+ = +1$ and $V_- = -1$ respectively, for times $s = [0; 0.3; 1.2]$. The initial conditions are given by $P_0(y) = Q_0(y) = \frac{1}{8\gamma} [\tanh(y + \gamma) - \tanh(y - \gamma)]$, $\alpha = 0$, $\beta = 2$ and $\gamma = \frac{1}{4}$.

In Fig. 4, the temporal evolution of the overall adoption rate $P(t) = \int_{\mathbb{R}} P(y, t) dy$ of the agents is illustrated and compared to the one observed for the Bass' original model. For the present configuration, the overall adoption rate stands between β , the rate observed for our spatial Bass' model in absence of imitation processes (Section 3.1.b), and $\beta + 1$, the rate obtained for the original aggregated Bass' model. Hence, in general, the overall adoption rate of our spatial Bass' model will be equal to $\beta + \epsilon(t)$, $\epsilon(t) \in]0, 1]$, with $\epsilon(t)$ depending on the initial distributions of the agents and on the model parameters. Remember that the imitation rate (and hence the overall adoption rate) is controlled by the number of *neighbors* that each agent effectively observes during the imitation process. In the aggregated (original) Bass' model, this number is maximum as each agent systematically observes the global population of agents. In the limiting regime of Section 3.1.b, the number of observed agents is equal to 0 as the two populations of agents \mathcal{A}_+ and \mathcal{A}_- are initially and hence




Figure 4: Temporal evolution of the overall adoption rate $P(t)$ *i)* for the original Bass' model, *ii)* for our spatial Bass' model when there are imitation processes (Section 3.1.c) and *iii)* for our spatial Bass' model when there are no imitation processes because the two populations of agents \mathcal{A}_+ and \mathcal{A}_- are immediately segregated (Section 3.1.b).

permanently segregated, thus allowing no imitation processes.

4 Conclusion and Perspectives

Often, agents' remoteness may naturally reduce the efficiency of imitation processes, a feature that is totally absent in the original aggregated Bass' approach. Incorporating a spatial dimension into the Bass' dynamics is however not a minor extension. It transforms indeed a nonlinear single dimension dynamics into a nonlinear infinite dimensional field dynamics for which, in general, no solution methods are available. It is hence truly remarkable that our spatial generalization of the original Bass' dynamics leads to a class of models for which the evolution of the resulting spatio-temporal patterns can be completely and exactly calculated. Indeed, the quadratic nonlinearity, due here to the underlying imitation mechanism, coincides with the collision term found in a solvable Boltzmann equation, the RW dynamics, used for gas models in mathematical physics. Our paper points out that the exceptional beauty of the RW dynamics, solvable via a linearizing logarithmic transformation, offers a unique, synthetic

and exact modeling framework to study nonlinear features generated by spatio-temporal imitation processes in economics systems.

References

- [1] E. Mansfield. *Technical Change and the Rate of Imitation*. *Econometrica*, **29**:741-766, (1961).
- [2] Z. V. Griliches. *Hybrid Corn: An Exploration in the Economics of Technological Change*. *Econometrica*, **25**:501-522, (1957).
- [3] Z. V. Griliches. *Hybrid Corn and the Economics of Innovation*. *Science*, **132**:275-280, (1960).
- [4] F. Bass. *A New Product Growth Model for Consumer Durables*. *Management Science*, **15**(5):215-227, (1969).
- [5] J. A. Schumpeter. *Business Cycles: A Theoretical, Historical and Statistical Analysis of the Capitalist Process*. McGraw-Hill, New York, (1939).
- [6] B. Jovanovic and R. Rob. *Long Waves and Short Waves: Growth Through Intensive and Extensive Search*. *Econometrica*, **58**(6):1391-1409, (1990).
- [7] R. Pastor-Satorras and A. Vespigniani. *Epidemic Spreading in Scale-Free Networks*. *Physics Review Letters*, **86**:3200-3203, (2001).
- [8] E. Glaeser, B. Sacerdote and J. Scheinkman. *Crime and Social Interactions*. *Quarterly Journal of Economics*, **111**:507-548, (1996).
- [9] R. Cont and J. P. Bouchard. *Herd Behavior and Aggregate Fluctuations in Financial Market*. *Macroeconomic Dynamics*, **4**(2):170-196, (2000).
- [10] A. Corcos, J.-P. Eckmann, A. Malaspinas, Y. Malevergne and D. Sornette. *Imitation and Contrarian Behavior: Hyperbolic Bubbles, Crashes and Chaos*. *Quantitative Finance*, **2**:264-281, (2002).
- [11] P. Dai Pra, W. J. Runggaldier, E. Sartori and M. Tolotti. *Large Portfolio Losses: A Dynamic Contagion Model*. *Annals of Applied Probability*, **19**(1):347-394, (2009).
- [12] U. Horst. *Stochastic Cascades, Credit Contagion and Large Portfolio Losses*. *Journal of Economic Behavior & Organization*, **63**:25-54, (2007).
- [13] D. Lopez-Pintado and D. J. Watts. *Social Influence, Binary Decisions and Collective Dynamics*. *Rationality and Society* **20**(4):399-443, (2008).

- [14] D. Lopez-Pintado. *Diffusion in Complex Social Networks*. Games and Economic Behavior, **62**:573-590, (2008).
- [15] F. Collet, P. Dai Pra and E. Sartori. *A Simple Mean-Field Model for Social Interactions: Dynamics, Fluctuations, Criticality*. Journal of Statistical Physics, **139**:820-858, (2010).
- [16] T. W. Valente. *Network Models of Diffusion of Innovations*. Hampton Press ,(1995).
- [17] W. Brock and S. N. Durlauf. *Discrete Choice with Social Interactions*. Review of Economic Studies, **68**:235-260, (2001).
- [18] L. Blume. *The Statistical Mechanics of Strategic Interactions*. Games and Economic Theory, **5**:387-424, (1993).
- [19] M. Schultz. *Statistical Physics and Economics - Concepts, Tools and Applications*. Springer Verlag, (2003).
- [20] L. Blume and S. N. Durlauf. *The Interactions-Based Approach to Socioeconomic Behavior*. Mimeo, Department of Economics, University of Wisconsin, (1998).
- [21] W. Brock and S. N. Durlauf. *Interaction-Based Model*. In J. Heckmann and E. Leamer (Eds.), Handbook of Econometrics, **5**:3297-3380, North Holland, (2001).
- [22] G. Ellison. *Learning, Local Interactions and Coordination*. Econometrica, **61**(5):1047-1071, (1993).
- [23] G. Ellison and D. Fudenberg. *Rules of Thumb for Social Learning*, Journal of Political Economy, **101**:612644, (1993).
- [24] R. Andergassen, F. Nardini and M. Ricottilli. *Innovation Waves, Self-Organized Criticality and Technological Convergence*. Journal of Economic Behavior & Organization, **61**:710728, (2006).
- [25] D. Levine. *Is Behavioral Economics Doomed?*. Max Weber Lecture, (June 2009).
- [26] D. Levine and D. Fudenberg. *Learning and Equilibrium*. Annual Review of Economics, **1**(1): 385-419, (2008).
- [27] D. Levine and W. Pesendorfer. *Evolution of Cooperation Through Imitation*. Games and Economic Behavior, **58**: 293-315, (2007).
- [28] U. Horst. *Dynamic Systems of Social Interactions*. Journal of Economic Behavior & Organization, **73**:158-170, (2010).
- [29] A. Foster and M. Rosenzweig. *Learning by Doing and Learning from Others: Human Capital and Technical Change in Agriculture*. Journal of Political Economy, **103**:1176-1209, (1995).

- [30] E. M. Rogers. *Diffusion of Innovations*. Free Press, New York, (1992).
- [31] P. Krugman. *Geography and Trade*. MIT Press, Cambridge, MA, (1991).
- [32] M. P. Feldman. *Knowledge Complementary and Innovation*. Small Business Economics, **6**:363-372, (1994).
- [33] M. P. Feldman. *The Geography of Innovation*. Kluwer Academic Publishers, Boston, (1994).
- [34] A. C. Case. *Spatial Patterns in Household Demand*. *Econometrica*, **59**(4):953-965,(1991).
- [35] D. B. Audretsch and M. P. Feldman. *Knowledge Spillovers and the Geography of Innovation and Production*. *American Economic Review*, **86**:630-640, (1966).
- [36] Z. J. Acs, D. B. Audretsch and M. P. Feldman. *Real Effects of Academic Research: Comment*. *American Economic Review*, **82**(1): 363-367, (1992).
- [37] Z. J. Acs, D. B. Audretsch and M. P. Feldman. *R&D Spillovers and Recipient Firm Size*. *Review of Economics and Statistics*, **76**: 336-340, (1994).
- [38] A. Jaffe, M. Trajtenberg and R. Henderson. *Geographic Localisation of Knowledge Spillovers as Evidenced by Patent Citations*. *Quarterly Journal of Economics*, **108**:577-598, (1993).
- [39] G. A. Akerlof. *Social Distance and Social Decisions*. *Econometrica* **65**(5):1005-1027, (1997).
- [40] T. Vicsek, A. Czirok, E. Ben-Jacob, I. Cohen and O. Shochet. *Novel Type of Phase Transition in a System of Self-Driven Particles*. *Physical Review Letters*, **75**:12261-229, (1995).
- [41] F. Cucker and S. Smale. *Emergent Behavior in Flocks*. *IEEE Transactions on Automatic Control*, **52**(5):852-862, (2007).
- [42] T. W. Ruikgrok and T. T. Wu. *A Completely Solvable Model of the Nonlinear Boltzmann Equation*. *Physica A*, **113**:401-416, (1982).
- [43] M.-O. Hongler, O. Gallay, M. Hülsmann, P. Cordes and R. Colmorn. *Centralized Versus Decentralized Control - A Solvable Stylized Model in Transportation*. *Physica A*, **389**:4162-4171, (2010).
- [44] M.-O. Hongler and L. Streit. *A Probabilistic Connection Between the Burgers and a Discrete Boltzmann Equation*. *Europhysics Letters*, **12**:193-197, (1990).

- [45] V. Mahajan and R. A. Peterson. *Integrating Time and Space in Technological Substitution Models*. Technological Forecasting Social Change, **14**:231-241, (1979).
- [46] W. M. Cohen and D. A. Levinthal. *Innovation and Learning: The Two Faces of R and D*. Economic Journal, **99**:569-596, (1989).
- [47] E. Gutkin and M. Kac. *Propagation of Chaos and the Burgers' Equation*. SIAM Journal of Applied Mathematics, **434**(4):971-980, (1963).
- [48] M. Abramowitz and I. Stegun. *Handbook of Mathematical Functions*. Dover, see entry 9.6.