
Variance reduction

Michel Bierlaire

`michel.bierlaire@epfl.ch`

Transport and Mobility Laboratory

Example

- Use simulation to compute

$$I = \int_0^1 e^x dx$$

- We know the solution: $e - 1 = 1.7183$
- Simulation: consider draws two by two
- Let r_1, \dots, r_R be independent draws from $U(0, 1)$.
- Let s_1, \dots, s_R be independent draws from $U(0, 1)$.

$$I \approx \frac{1}{R} \sum_{i=1}^R \frac{e^{r_i} + e^{s_i}}{2}$$

- Use $R = 10'000$ (that is, a total of 20'000 draws)
- Mean over R draws from $(e^{r_i} + e^{s_i})/2$: 1.720, variance: 0.123


Example

- Now, use half the number of draws
- Idea: if $X \sim U(0, 1)$, then $(1 - X) \sim U(0, 1)$
- Let r_1, \dots, r_R be independent draws from $U(0, 1)$.

$$I \approx \frac{1}{R} \sum_{i=1}^R \frac{e^{r_i} + e^{1-r_i}}{2}$$

- Use $R = 10'000$
- Mean over R draws of $(e^{r_i} + e^{1-r_i})/2$: 1.7183, variance: **0.00388**
- Compared to: mean of $(e^{r_i} + e^{s_i})/2$: 1.720, variance: **0.123**

Example



Antithetic draws

- Let X_1 and X_2 i.i.d r.v. with mean θ
- Then

$$\text{Var} \left(\frac{X_1 + X_2}{2} \right) = \frac{1}{4} (\text{Var}(X_1) + \text{Var}(X_2) + 2 \text{Cov}(X_1, X_2))$$

- If X_1 and X_2 are independent, then $\text{Cov}(X_1, X_2) = 0$.
- If X_1 and X_2 are negatively correlated, then $\text{Cov}(X_1, X_2) < 0$, and the variance is reduced.

Back to the example

- Independent draws
- $X_1 = e^U, X_2 = e^U$

$$\begin{aligned}\text{Var}(X_1) = \text{Var}(X_2) &= \mathbb{E}[e^{2U}] - \mathbb{E}[e^U]^2 \\ &= \int_0^1 e^{2x} dx - (e - 1)^2 \\ &= \frac{e^2 - 1}{2} - (e - 1)^2 \\ &= 0.2420\end{aligned}$$

$$\text{Cov}(X_1, X_2) = 0$$

$$\text{Var}\left(\frac{X_1 + X_2}{2}\right) = \frac{1}{4} (0.2420 + 0.2420) = 0.1210$$

Back to the example

- Antithetic draws
- $X_1 = e^U, X_2 = e^{1-U}$

$$\text{Var}(X_1) = \text{Var}(X_2) = 0.2420$$

$$\begin{aligned}\text{Cov}(X_1, X_2) &= E[e^U e^{1-U}] - E[e^U]E[e^{1-U}] \\ &= e - (e-1)(e-1) \\ &= -0.2342\end{aligned}$$

$$\text{Var}\left(\frac{X_1 + X_2}{2}\right) = \frac{1}{4} (0.2420 + 0.2420 - 2 \cdot 0.2342) = 0.0039$$

Antithetic draws: generalization

- Suppose that

$$X_1 = h(U_1, \dots, U_m)$$

where U_1, \dots, U_m are i.i.d. $U(0, 1)$.

- Define

$$X_2 = h(1 - U_1, \dots, 1 - U_m)$$

- X_2 has the same distribution as X_1
- If h is monotone in each of its coordinates, then X_1 and X_2 are negatively correlated.

Control variates

- We use simulation to estimate $\theta = E[X]$, where X is an output of the simulation
- Let Y be another output of the simulation, such that we know $E[Y] = \mu$
- We consider the quantity:

$$Z = X + c(Y - \mu).$$

- By construction, $E[Z] = E[X]$
- Its variance is

$$\text{Var}(Z) = \text{Var}(X + cY) = \text{Var}(X) + c^2 \text{Var}(Y) + 2c \text{Cov}(X, Y)$$

- Find c such that $\text{Var}(Z)$ is minimum

Control variates

- First derivative:

$$2c \operatorname{Var}(Y) + 2 \operatorname{Cov}(X, Y)$$

- Zero if

$$c^* = -\frac{\operatorname{Cov}(X, Y)}{\operatorname{Var}(Y)}$$

- Second derivative:

$$2 \operatorname{Var}(Y) > 0$$

- We use

$$Z^* = X - \frac{\operatorname{Cov}(X, Y)}{\operatorname{Var}(Y)}(Y - \mu).$$

- Its variance

$$\operatorname{Var}(Z^*) = \operatorname{Var}(X) - \frac{\operatorname{Cov}(X, Y)^2}{\operatorname{Var}(Y)} \leq \operatorname{Var}(X)$$

Control variates

In practice...

- $\text{Cov}(X, Y)$ and $\text{Var}(Y)$ are usually not known.
- We can use their sample estimates:

$$\widehat{\text{Cov}}(X, Y) = \frac{1}{n-1} \sum_{r=1}^R (X_r - \bar{X})(Y_r - \bar{Y})$$

and

$$\widehat{\text{Var}}(Y) = \frac{1}{n-1} \sum_{r=1}^R (Y_r - \bar{Y})^2.$$

Control variates

In practice...

- Alternatively, use linear regression

$$X = aY + b + \varepsilon$$

where $\varepsilon \sim N(0, \sigma^2)$.

- The least square estimators of a and b are

$$\hat{a} = \frac{\sum_{r=1}^R (X_r - \bar{X})(Y_r - \bar{Y})}{\sum_{r=1}^R (Y_r - \bar{Y})^2}$$

$$\hat{b} = \bar{X} - \hat{a}\bar{Y}.$$

- Therefore

$$c^* = -\hat{a}$$

Control variates

- Moreover,

$$\begin{aligned}\hat{b} + \hat{a}\mu &= \bar{X} - \hat{a}\bar{Y} + \hat{a}\mu \\ &= \bar{X} - \hat{a}(\bar{Y} - \mu) \\ &= \bar{X} + c^*(\bar{Y} - \mu) \\ &= \hat{\theta}\end{aligned}$$

- Therefore, the control variate estimate $\hat{\theta}$ of θ is obtained by the estimated linear model, evaluated at μ .


Back to the example

- Use simulation to compute $I = \int_0^1 e^x dx$
- $X = e^U$
- $Y = U$, $E[Y] = 1/2$, $\text{Var}(Y) = 1/12$
- $\text{Cov}(X, Y) = (3 - e)/2 \approx 0.14$
- Therefore, the best c is

$$c^* = -\frac{\text{Cov}(X, Y)}{\text{Var}(Y)} = -6(3 - e) \approx -1.69$$

- Test with $R = 10'000$
- Result of the regression: $\hat{a} = 1.6893$, $\hat{b} = 0.8734$
- Estimate: $\hat{b} + \hat{a}/2 = 1.7180$, Variance: 0.003847 (compared to 0.24)

Back to the example



Variance reductions techniques

- Conditioning
- Stratified sampling
- Importance sampling
- Draw recycling

In general: correlation helps!