
Simulating events: the Poisson process

Michel Bierlaire

`michel.bierlaire@epfl.ch`

Transport and Mobility Laboratory

Siméon Denis Poisson



Siméon-Denis Poisson (1781–1840). French mathematician.

- Poisson random variable
- Poisson process
- Non homogeneous Poisson process

Poisson random variable


- Number of successes in a large number n of trials (binomial distribution)
- when the probability p of a success is small.
- Denote $\lambda = np$.

$$\Pr(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}.$$


Property:

$$E[X] = \text{Var}(X) = \lambda.$$

Poisson random variable



Poisson random variable



Poisson process

- Events are occurring at random time points
- $N(t)$ is the number of events during $[0, t]$
- They constitute a Poisson process with rate $\lambda > 0$ if
 1. $N(0) = 0$,
 2. # of events occurring in disjoint time intervals are independent,
 3. distribution of $N(t + s) - N(t)$ depends on s , not on t ,
 4. probability of one event in a small interval is approx. λh :

$$\lim_{h \rightarrow 0} \frac{\Pr(N(h) = 1)}{h} = \lambda,$$

5. probability of two events in a small interval is approx. 0:

$$\lim_{h \rightarrow 0} \frac{\Pr(N(h) \geq 2)}{h} = 0.$$

Poisson process

Property:

$$N(t) \sim \text{Poisson}(\lambda t), \quad \Pr(N(t) = k) = e^{-\lambda t} \frac{(\lambda t)^k}{k!}$$

Inter-arrival times:

- S_k is the time when the k th event occurs,
- $X_k = S_k - S_{k-1}$ is the time elapsed between event $k - 1$ and event k .
- $X_1 = S_1$
- Distribution of X_1 : $\Pr(X_1 > t) = \Pr(N(t) = 0) = e^{-\lambda t}$.
- Distribution of X_2 :

$$\begin{aligned} \Pr(X_k > t | S_{k-1} = s) &= \Pr(0 \text{ events in }]s, s + t] | S_{k-1} = s) \\ &= \Pr(0 \text{ events in }]s, s + t]) \\ &= e^{-\lambda t}. \end{aligned}$$

Poisson process

- X_1 is an exponential random variable with mean $1/\lambda$
- X_2 is an exponential random variable with mean $1/\lambda$
- X_2 is independent of X_1 .
- Same arguments can be used for $k = 3, 4, \dots$

Therefore, the CDF of X_k is, for any k ,

$$F(t) = \Pr(X_k \leq t) = 1 - \Pr(X_k > t) = 1 - e^{-\lambda t}.$$

The pdf is

$$f(t) = \frac{dF(t)}{dt} = \lambda e^{-\lambda t}.$$

The inter-arrival times X_1, X_2, \dots are independent and identically distributed exponential random variables with parameter λ , and mean $1/\lambda$.

Poisson process

- Simulation of event times of a Poisson process with rate λ until time T :
 1. $t = 0, k = 0$.
 2. Draw $r \sim U(0, 1)$.
 3. $t = t - \ln(r)/\lambda$.
 4. If $t > T$, STOP.
 5. $k = k + 1, X_k = t$.
 6. Go to step 2.

Non homogeneous Poisson process

- Assume that the rate varies with time, and call it $\lambda(t)$.
- The events constitute a non homogeneous Poisson process with rate $\lambda(t)$ if
 1. $N(0) = 0$
 2. # of events occurring in disjoint time intervals are independent,
 3. probability of one event in a small interval is approx. $\lambda(t)h$:

$$\lim_{h \rightarrow 0} \frac{\Pr((N(t+h) - N(t)) = 1)}{h} = \lambda(t),$$

4. probability of two events in a small interval is approx. 0:

$$\lim_{h \rightarrow 0} \frac{\Pr((N(t+h) - N(t)) \geq 2)}{h} = 0.$$

Non homogeneous Poisson process

- Mean value function:

$$m(t) = \int_0^t \lambda(s) ds, \quad t \geq 0.$$

- Poisson distribution:

$$N(t + s) - N(t) \sim \text{Poisson}(m(t + s) - m(t))$$

- Link with homogeneous Poisson process:
 - Consider a Poisson process with rate λ .
 - If an event occurs at time t , count it with probability $p(t)$.
 - The process of counted events is a non homogeneous Poisson process with rate $\lambda(t) = \lambda p(t)$.

Non homogeneous Poisson process

Proof:

1. $N(0) = 0$ [OK]
2. # of events occurring in disjoint time intervals are independent, [OK]
3. probability of one event in a small interval is approx. $\lambda(t)h$: [?]

$$\lim_{h \rightarrow 0} \frac{\Pr((N(t+h) - N(t)) = 1)}{h} = \lambda(t),$$

4. probability of two events in a small interval is approx. 0: [OK]

$$\lim_{h \rightarrow 0} \frac{\Pr((N(t+h) - N(t)) \geq 2)}{h} = 0.$$

Non homogeneous Poisson process

- $N(t)$ number of events of the non homogeneous process
- $N'(t)$ number of events of the underlying homogeneous process

$$\begin{aligned}\Pr((N(t+h) - N(t)) = 1) &= \sum_k \Pr((N'(t+h) - N'(t)) = k, 1 \text{ is counted}) \\ &= \Pr((N'(t+h) - N'(t)) = 1, 1 \text{ is counted}) \\ &= \Pr((N'(t+h) - N'(t)) = 1) \Pr(1 \text{ is counted}) \\ &= \Pr(N'(h) = 1) \Pr(1 \text{ is counted})\end{aligned}$$

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{\Pr((N(t+h) - N(t)) = 1)}{h} &= \lim_{h \rightarrow 0} \frac{\Pr(N'(h) = 1)}{h} \Pr(1 \text{ is counted}) \\ &= \lambda p(t).\end{aligned}$$

Non homogeneous Poisson process

Simulation of event times of a non homogeneous Poisson process with rate $\lambda(t)$ until time T :

1. Consider λ such that $\lambda(t) \leq \lambda$, for all $t \leq T$.
2. $t = 0$, $k = 0$.
3. Draw $r \sim U(0, 1)$.
4. $t = t - \ln(r)/\lambda$.
5. If $t > T$, STOP.
6. Generate $s \sim U(0, 1)$.
7. If $s \leq \lambda(t)/\lambda$, then $k = k + 1$, $X(k) = t$.
8. Go to step 3.

Summary

- Poisson random variable
- Poisson process
- Non homogeneous Poisson process
- Main assumption: events occur continuously and independently of one another
- Typical usage: arrivals of customers in a queue
- Easy to simulate